

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

September 1992

Book Review: *Revision of the genus *Spermophagus* Schoenherr (Coleoptera, Bruchidae, Amblycerinae)*

John M. Kingsolver

Florida State Collection of Arthropods, Florida Department of Agriculture and Consumer Services,
Gainesville, FL

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>

 Part of the [Entomology Commons](#)

Kingsolver, John M., "Book Review: *Revision of the genus *Spermophagus* Schoenherr (Coleoptera, Bruchidae, Amblycerinae)* " (1992). *Insecta Mundi*. 439.

<https://digitalcommons.unl.edu/insectamundi/439>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Book Review

Revision of the genus *Spermophagus* Schoenherr (Coleoptera, Bruchidae, Amblycerinae). Lech Borowiec. Genus, International Journal of Invertebrate Taxonomy (Supplement), Biologica Silesiae. 1991. 198 p. (Price not given).

Species in the genus *Spermophagus* are found in all of the Old World regions except Australia. Prior to this fine monograph, the taxonomy was a hodgepodge of local faunal treatments and single species descriptions with few usable keys or illustrations. Now we have available keys to 90 species with full descriptions, geographical distributions, and excellent illustrations of dorsal habitus, male genitalia, and many of the female genital valves.

Host plants of species of *Spermophagus* are to be found mostly in the Convolvulaceae and Malvaceae, but isolated species have been associated with Asteraceae, Caesalpiniaceae, Fabaceae, and Mimosaceae. Host lists include a number of economically important weedy plant species, especially in the Convolvulaceae, and might lead to a consideration of the bruchids as biocontrol agents.

The author is a member of the faculty at the University of Wroclaw in Wroclaw, Poland. - **John M. Kingsolver, c/o Florida State Collections of Arthropods, P.O. Box 147100, Gainesville, FL 32614-7100.**